

Neonatal Pain Assessment and Procedural Pain Management

Version: 5

This is a CONTROLLED document for internal use only, valid only if accessed from the Policies and Procedures site.

Original approval by NICU Interprofessional Practice Nov 26, 2010

Content

- 1.0 [Introduction](#)
- 2.0 [Definitions](#)
- 3.0 [Clinical Practice Recommendations](#)
 - 3.1 [Pain Assessment](#)
 - 3.1.1 [Pain assessment scores](#)
 - 3.1.2 [Frequency of pain assessment scores](#)
 - 3.2 [Pain Management](#)
 - 3.2.1 [Developmental strategies](#)
 - 3.2.2 [Topical lidocaine and prilocaine analgesia](#)
 - 3.2.3 [Sucrose administration](#)
 - 3.3 [Adjunct Procedural Pain Management](#)
 - 3.4 [Procedural Pain Management](#) (e.g., PICC, lumbar puncture, eye exams, chest tube, injections)
- 4.0 [Related Documents](#)
- 5.0 [References](#)

1.0 Introduction

- Pain management strategies will be focused on managing pain during procedures and managing surgical and subacute pain or disease related pain.
- In addition to developmental strategies, pharmacological strategies should be used based on the type of procedure.
- Refer to [Neonatal Sedation Management](#) guideline for additional guidance as neonate's may also be concomitantly receiving adjunct therapy with sedation medications.

2.0 Definitions

- **PIPP-R:** Premature Infant Pain Profile is a bio-behavioural observational tool for acute and procedural pain.
- **FLACC-R:** Face, Legs, Activity, Cry, Consolability is a behavioural observational tool for acute pain.

©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.

3.0 Clinical Practice Recommendations

The grading system in [Table 1](#) serves as a guideline for the user about the hierarchy of evidence available to support each recommendation.

Table 1: Grades of Recommendation	
A	Recommendation supported by at least one randomized controlled trial, systematic review or meta-analysis.
B	Recommendation supported by at least one cohort comparison, case study or other experimental study.
C	Recommendation supported by expert opinion or experience of a consensus panel.

3.1 Pain Assessment

3.1.1 Pain assessment scores

1. **PIPP-R score:** Premature Infant Pain Profile-Revised is a bio-behavioural observational tool for acute and procedural pain. PIPP-R scores to be completed for infants ≤ 48 weeks post menstrual age (i.e., 2 months corrected age).
2. **r-FLACC score:** revised Face, Legs, Activity, Cry, Consolability (*FLACC*) score is a behavioural observational tool for acute pain. r-FLACC scores are to be completed for infants > 48 weeks post menstrual age.

Example:

- 8-week-old 32-week gestation infant = 40 weeks gestation, therefore complete **PIPP-R** score.
 - 8-week-old 41-week gestation infant = 49 weeks gestation, therefore complete **r-FLACC** score.
- See [Pain Assessment](#).

3.1.2 Frequency of pain assessment scores

1. **Once per shift**, a pain score (PIPP-R or r-FLACC) scores to be assessed and documented on each patient, on admission, and before, during and after an invasive procedure as per hospital policy.
2. Pain scores are to be done more frequently for the following infants.
 - To assess pain in infants with known medical conditions or interventions that may cause pain (e.g., NEC, chest tubes).
 - Following post-operative procedures as per See [Pain Management Guidelines for Post-Operative Neonates](#)
 - Following changes to continuous analgesia.
 - To evaluate for pharmacological treatments for infants on short term opioids. For patients with treatment lengths greater than 5 days utilize routine Neonatal Abstinence Scoring for withdrawal of opioids as per the [Prevention and Treatment of Opioid and Benzodiazepine Withdrawal](#).

3.2 Pain Management

For general principles of pain management, see [Pain Management Guideline](#).

3.2.1 Developmental strategies

- All infants should receive physical/psychological developmentally appropriate strategies during all painful procedures² **(Grade B)**.
- Developmental strategies that are considered acceptable for tissue damaging procedures (such as needle pokes) include:
 1. Sucrose
 2. Skin to skin contact
 3. Breastfeeding
- Other developmental strategies are considered adjunct strategies. They can be combined with one of the above strategies or with pharmacological strategies to reduce pain during procedures. These strategies include:
 1. Non-nutritive sucking
 2. Positioning and containment
 3. Swaddling
 4. Reduction of light and sound levels
 5. Minimal handling
 6. Auditory and visual distraction

3.2.2 Topical lidocaine and prilocaine analgesia

- Local topical analgesics may be used with some procedures but may be limited if vasoconstriction may inhibit the success of the procedure (e.g., IV insertion). **(Grade C)**.
- Topical Lidocaine and Prilocaine analgesia (e.g., EMLA) (see [SickKids e-formulary](#))⁸ can be used in all infants.
- If infant is <1000 grams or < 14 days old, consider using gauze and avoiding use of an occlusive dressing (e.g., Tegaderm™) to prevent skin injury during removal.


3.2.3 Sucrose administration

- **Refer to [SickKids e-Formulary](#) for dosing of sucrose in neonates.** Doses may be repeated for prolonged procedures.
- Document sucrose uses and effectiveness using appropriate pain scores.
- Order sucrose on a PRN basis to facilitate use for procedures such as bloodwork, IV starts, eye exams, drain removal, echocardiography, nasogastric or orogastric tube insertion.
- Sucrose may be ordered by the MD/NP or RN ([under Nursing Order Policy](#)).
- Refer to [SickKids e-Formulary](#) for sucrose contraindications
- Refer to [Pain Management Guideline](#).

3.3 Adjunct Procedural Pain Management

The following recommended pain management strategies are based on the estimated severity of pain. In addition, use appropriate developmental pain reduction strategies as outlined above.

Estimated Pain Severity of NICU Procedures (Laudiano-Dray et al. 2020)



©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.

3.4 Procedural Pain Management: Recommended pain management options for neonatal procedures.

Procedure	Recommended Pain Reduction Management
Chest tube insertion	<ul style="list-style-type: none"> - Fentanyl 1mcg/kg/dose IV 3 - 5 minutes prior to procedure. - Use pacifier with 24% Sucrose 2 minutes prior to procedure SickKids e-formulary. - Lidocaine 1% subcutaneously (SQ) as local anesthetic as per SickKids e-formulary. - Start morphine infusion of 5 mcg/kg/hr following opioid bolus and assess infant pain scores. - PRN morphine and/or IV or enteral acetaminophen can also be used.
Chest tube removal	<ul style="list-style-type: none"> - Fentanyl bolus 0.5mcg/kg/dose IV 3-5 minutes prior to procedure. - Use pacifier with 24% Sucrose 2 minutes prior to procedure as per SickKids e-formulary (Grade B).
Echocardiogram	<ul style="list-style-type: none"> - Use pacifier with 24% sucrose 2 minutes prior to procedure as per sucrose administration guideline and SickKids e-formulary (Grade B).
Eye exams (e.g., ROP exam)	<ul style="list-style-type: none"> - Use pacifier with 24% Sucrose 2 minutes prior to procedure SickKids e-formulary (Grade B) in addition to eye drops prescribed by Ophthalmology.
Eye - intravitreal bevacizumab (i.e., Avastin)	<ul style="list-style-type: none"> - Administer Fentanyl 0.5-1 mcg/kg/dose IV 3 - 5 minutes prior to the procedure. - Consider using midazolam 0.05 - 0.1 mg/kg/dose IV prn as per SickKids e-formulary for sedation. Use cautiously in non-intubated infants. - Use pacifier with 24% Sucrose 2 minutes prior to procedure SickKids e-formulary (Grade B). - Use developmental strategies such as bundling for containment during the procedure.
Heel lance	<ul style="list-style-type: none"> - Use pacifier with 24% Sucrose 2 minutes prior to procedure as per SickKids e-Formulary (Grade B), skin to skin or breastfeeding.
Injection - intramuscular (e.g., immunization)	<ul style="list-style-type: none"> - Use pacifier with 24% Sucrose 2 minutes prior to procedure as per SickKids e-Formulary (Grade B), skin to skin or breastfeeding. - Apply topical anesthetic cream EMLA (lidocaine and prilocaine) 45 – 60 minutes prior to procedure as per SickKids e-Formulary. - Complete injection as per Intramuscular Injections policy and procedure.
Injection - subcutaneous	<ul style="list-style-type: none"> - Use pacifier with 24% Sucrose 2 minutes prior to procedure as per SickKids e-formulary (Grade B), skin to skin or breastfeeding. - Complete injection as per Intramuscular Injections policy and procedure.
Intubation	<ul style="list-style-type: none"> - As per Neonatal Endotracheal Intubation (Rapid Sequence Intubation).
Lumbar puncture	<ul style="list-style-type: none"> - Apply local topical anesthetic cream- EMLA 45-60 minutes prior to procedure as per SickKids e-formulary. - Consider fentanyl 1 mcg/kg/dose IV prior to procedure and/or midazolam 0.05 - 0.1 mg/kg/dose IV for sedation if infant is difficult to position. <ul style="list-style-type: none"> - Use midazolam cautiously in non-intubated infants. - For preterm infants < 35 weeks gestation, midazolam is contraindicated. - Use pacifier with 24% Sucrose 2 minutes prior to procedure SickKids e-formulary (Grade B). - Cautious physical handling is advised.

©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.

Procedure	Recommended Pain Reduction Management
Nasogastric or orogastric tube insertion	<ul style="list-style-type: none"> - Use pacifier with 24% sucrose 2 minutes prior to procedure SickKids e-formulary (Grade B).
Palliative care	<ul style="list-style-type: none"> - Physical and psychological strategies for pain management. - Oral, sublingual, or buccal morphine and/or lorazepam may be utilized as recommended by the palliative care team. - Refer to NICU End of life (EOL) guidelines and EOL order set.
PICC insertion	<ul style="list-style-type: none"> - Refer to table 2.
PICC removal for cuffed IGT lines only	<ul style="list-style-type: none"> - Cuffed Image Guided Therapy (IGT) lines must be removed by IGT staff and will be organized by Vascular access service (VAS) staff. - Procedure may be performed in NICU or IGT as determined by VAS and IGT. - Analgesia is as per vascular access service (VAS) and IGT recommendations. - Topical EMLA is often used as per SickKids e-formulary. - Uncuffed lines may be removed by NICU staff at the discretion of IGT.
Peripheral arterial sampling and peripheral arterial catheter insertion	<ul style="list-style-type: none"> - Apply topical anesthetic cream EMLA 45 – 60 minutes prior to procedure as per SickKids e-formulary. - If topical EMLA (i.e., lidocaine and prilocaine) is not used, fentanyl 1 mcg/kg/dose IV 3 - 5 minutes prior to procedure. - Ultrasound guided insertion should be used if available. Use of ultrasound may decrease analgesic requirements. - Use pacifier with 24% sucrose 2 minutes prior to the procedure as per SickKids e-formulary.
Post-operative pain management	<ul style="list-style-type: none"> - Refer to Neonatal Post-Operative Pain Guidelines.
Umbilical catheter insertion	<ul style="list-style-type: none"> - Use pacifier with 24% sucrose 2 minutes prior to procedure as per SickKids e-formulary.
Urinary catheter insertion Suprapubic bladder tap	<ul style="list-style-type: none"> - Use pacifier with 24% sucrose 2 minutes prior to procedure as per SickKids e-formulary.
Venipuncture or intravenous catheter insertion	<ul style="list-style-type: none"> - Use pacifier with 24% sucrose 2 minutes prior to procedure) as per SickKids e-formulary. - If desired and non-urgent, consider applying local topical anesthetic cream- EMLA (i.e., lidocaine and prilocaine) 45 - 60 minutes prior to procedure as per SickKids e-formulary.

Table 2: Peripherally Inserted Central Catheter (PICC) Recommended Pain Reduction Management			
PICC insertion - NICU	Preterm infants	Non-intubated <ul style="list-style-type: none"> - Fentanyl 0.5 mcg/kg/dose IV 3 – 5 minutes prior to procedure. - Use pacifier with 24% sucrose 2 minutes prior to procedure as per SickKids e-formulary (Grade B). 	Intubated <ul style="list-style-type: none"> - Fentanyl 1 mcg/kg/dose IV 3 – 5 minutes prior to procedure. - Use pacifier with 24% sucrose 2 minutes prior to procedure as per SickKids e-formulary (Grade B).
	Term infants	Intubated or non-intubated <ul style="list-style-type: none"> - Fentanyl 1 mcg/kg/dose IV 3 – 5 minutes prior to procedure. - Administer fentanyl by slow IV over 3 - 5 minutes OR Morphine 0.1 mg/kg/dose IV 20 minutes prior to procedure. - Consider using midazolam 0.05-0.1mg/kg/dose prn as per SickKids e-formulary if needed for sedation. Use cautiously in non-intubated infants. - Use pacifier with 24% sucrose 2 minutes prior to procedure as per sucrose administration guideline and SickKids e-formulary (Grade B). 	
PICC insertion – IGT (Image guided therapy)	Preterm infants	Non-intubated <ul style="list-style-type: none"> - Fentanyl 0.5 mcg/kg/dose IV 3 – 5 minutes prior to procedure. - Use pacifier with 24% sucrose 2 minutes prior to procedure as per SickKids e-Formulary (Grade B). - Apply topical anesthetic cream EMLA (lidocaine and prilocaine) 45 – 60 minutes prior to procedure as per SickKids e-formulary. 	Intubated <ul style="list-style-type: none"> - Fentanyl 1 mcg/kg/dose IV 3 – 5 minutes prior to procedure. - Use pacifier with 24% sucrose 2 minutes prior to procedure as per SickKids e-Formulary (Grade B). - Apply topical anesthetic cream EMLA (i.e., lidocaine and prilocaine). 45 – 60 minutes prior to procedure as per SickKids e-formulary.
	Term infants	Intubated or non-intubated <ul style="list-style-type: none"> - Fentanyl 1 mcg/kg/dose IV 3 – 5 minutes prior to procedure OR Morphine 0.1 mg/kg/dose IV 20 minutes prior to procedure. - Midazolam 0.05 mg/kg/dose IV 5 minutes prior to procedure if needed for additional sedation. - May repeat midazolam 0.05 mcg/kg/dose IV x 1, 30 minutes after first dose if inadequate sedation. - Use pacifier with 24% sucrose 2 minutes prior to procedure as per SickKids e-formulary (Grade B). - Apply topical anesthetic cream EMLA 45 – 60 minutes prior to procedure as per SickKids e-formulary. 	

4.0 Related Documents

[E-formulary: Lidocaine and Prilocaine \[EMLA \(eutectic mixture of local anesthetics\)\]](#)

[E-formulary: Sucrose](#)

[Neonatal Post-Operative Pain Guidelines](#)

[NICU Sedation Guidelines](#)

[Pain Assessment](#)

[Pain Management Guideline](#)

[Removal of a Peripherally Inserted Central Catheter \(PICC\)](#)

5.0 References

1. AAP Committee on Fetus and Newborn on Anesthesiology and Pain Medicine. (2016). Prevention and management of pain in the neonate: An update. *Archives of Pediatric Adolescent Medicine*, 137: e20154271.
2. Anand, KJS. (2007). Pharmacological approaches to the management of pain in the neonatal intensive care unit. *Journal of Perinatology*, 27(1), S4-11.
3. Bueno, M, Ballantyne, M, Campbell-Yeo, M, Estabrooks, CA, Gibbins, S, Harrison, D, McNair, C, Riahi, S, Squires, J, Synnes, A, Taddio, A, Victor, C, Yamada, J, Stevens, B. (2023) The effectiveness of repeated sucrose for procedural pain in neonates in longitudinal observational study. *Frontiers in Pain Research*, 4.
4. Bueno, M, Ballantyne, M, Campbell-Yeo, M, Estabrooks, C, Gibbins, S, Harrison, D, McNair, C, Riahi, S, Squires, J, Synnes, A, Taddio, A, Victor, C, Yamada, J, Stevens, B. (2024) Cumulative sucrose exposure for repeated procedural pain in preterm neonates and neurodevelopment at 18 months of corrected age: a prospective observational longitudinal study. *BMJ Paediatrics Open*, 8, e002604.
5. Campbell-Yeo, M, Disher, T, Mitra, S, Canadian Paediatric Society, Fetus and Newborn Committee. (2024). Managing pain in newborns: A multidimensional approach. Canadian Paediatric Society. URL: <https://cps.ca/en/documents/position/managing-pain-in-newborns>.
6. Doesburg, SM, Chau, CM, Cheung, TPL, Moiseev, A, Ribary, U, Herdman, AT, Miller, SP, Cepeda, IL, Synnes, A, Grunau, RE. (2104). Neonatal pain-related stress, functional cortical activity and visual-perceptual abilities in school-age children born at extremely low gestational age. *Pain*, 154(10):1946–52.
7. Franck, L.S. Lawhorn, G. (1998). Environmental and behavioural strategies to prevent and manage pain. *Seminars in Perinatology*, 22(5): 434-443.
8. Kasirer, Y, Shah, V, Yoon, EW, Bromiker, R, McNair, C, Taddio, A. (2018). Safety of fentanyl for peripherally inserted central catheter in non-intubated infants in the neonatal intensive care unit. *Journal of Perinatology*, 38(5), 526-529.
9. Khuran, S, Whit Hall, R, Anand, KJS. (2005). Treatment of pain and stress in the neonate: When and how. *Neoreviews*, 6(2): e76-e86.
10. Laudiano-Dray, MP, Pillai Riddell, R, Jones, L, Iyer, R, Whitehead, K, Fitzgerald, M, Fabrizi, L, Meek, J. (2020). Quantification of neonatal procedural pain severity: A platform for estimating total pain burden in individual infants. *Pain*, 161, 1270-1277.
11. McNair, C, Campbell-Yeo, M, Johnston, C, Taddio, A. (2019). Nonpharmacological management of pain during common needle puncture procedures in infants: Current research evidence and practical considerations: An update. *Clinics in Perinatology*, 46(4), 709-730.
12. Stevens, B., Yamada, J., Ohlsson, A. (2016). Sucrose for analgesia in newborn infants undergoing painful procedures. *Cochrane Database of Systemic Reviews*, 5.
13. Taddio A, Lee C, Yip A, Parvez B, McNamara PJ, Shah V. (2006). Intravenous morphine and topical tetracaine for treatment of pain in preterm neonates undergoing central line placement. *JAMA*, 295(7): 793-800.
14. Taddio, A., Ohlsson, A., Einarson, T., Stevens, B., Koren, G. (1998). A systematic review of lidocaine prilocaine (EMLA) in the treatment of acute pain in neonates. *Pediatrics*, 101(2), e1-9.
15. Vinall, J, Grunau, RE. (2014). Impact of repeated procedural pain-related stress in infants born very preterm. *Pediatric Research*, 75(5):584–7.

©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.

16. Walden, M. (2001) Pain Assessment and management: Guideline for practice. *National Association of Neonatal Nurses*, 1-24.
17. Walker, SM. (2019). Long-term effects of neonatal pain. *Semin Fetal Neonatal Med*, 24(4):101005.
18. Yamada, J, Bueno, M, Santos, L, Haliburton, S, Campbell-Yeo, M, Stevens, B. (2023) Sucrose analgesia for heel-lance procedures in neonates. *Cochrane Database of Systematic Reviews*, 8