

This is a CONTROLLED document for internal use only, valid only if accessed from the Policies and Procedures site.

### 1.0 Introduction

This pathway is for use with children aged 2-18 years old with no underlying disease or co-morbidity who have a confirmed diagnosis of appendicitis (either non-perforated or perforated) by the General Surgery Team and require a surgical appendectomy or medical management. Patients are to be removed from this pathway if there are significant postoperative complications (e.g. bowel obstruction or prolonged TPN) or a change in diagnosis.

The care pathways attached refer to patients admitted post-operatively. For patients with uncomplicated appendicitis discharged on day of surgery, discharge criteria are:

- Vital signs at baseline or within 10% of baseline vitals
- Good pain control: numeric rating score of less than 7/10 or pain word scale "medium" or less
- Tolerating clear fluids
- Able to transport safely home with family
- AVS reviewed with patient and family and PIV discontinued

### Target Users


- General Surgery Surgeons, Residents, Fellow, Nurse Practitioners, and nurses on the inpatient units

### 2.0 Definitions

- **Non-perforated appendicitis**- Appendix is normal, injected, inflamed or suppurative. Presence of cloudy fluid +/- local fibrinous exudate, or gangrenous appendicitis but with no visible hole in appendix, abscess or free fecalith
- **Perforated appendicitis**- Visible hole in appendix, free fecalith, diffuse peritonitis with fibrinous exudate or abscess
- **Appendectomy**- surgical removal of the appendix
- **Fever**- a fever is defined as any temperature reading greater than 38°C

### 3.0 Recommendations - [Printable Version](#)

**Appendicitis Algorithm**



\*If the patient clinically worsens, consider upgrade to IV piperacillin/tazobactam. If no ultrasound within the past 2-3 days, repeat in order to evaluate for drainable collection. If collection found, refer to algorithm for patient with drainable collections (IGT).

\*\*Alternative oral antibiotic therapy with Ciprofloxacin and Metronidazole may be considered in setting of confirmed beta-lactam allergy

\*\*\*Antibiotic therapy should be reassessed based on any available microbiological data (e- if cultures are obtained from an abdominal abscess aspiration)

\*\*\*\* A fever is defined as any temperature reading greater than 38° C (refer to Sepsis Clinical Pathway)

## Inpatient Non-Perforated Appendicitis Care Pathway

	PRE-OPERATIVE	POST-OPERATIVELY	DISCHARGE: WITHIN 24 HOURS POST-OP
GOALS	<ol style="list-style-type: none"> <li>1. Hydration maintained</li> <li>2. Adequate pain control</li> <li>3. Patient prepared for OR</li> <li>4. Child/family are advised of pre-op bath. Wipes to be used upon arrival. Refer to <a href="#">procedure document</a></li> </ol>	<ol style="list-style-type: none"> <li>1. Afebrile</li> <li>2. Adequate pain control</li> <li>3. Ambulating</li> <li>4. Able to tolerate diet (clear fluids to regular diet)</li> <li>5. Incision intact, no drainage, dry and intact</li> </ol>	<ol style="list-style-type: none"> <li>1. Afebrile</li> <li>2. Adequate pain control</li> <li>3. Ambulating</li> <li>4. Able to tolerate diet</li> <li>5. Incision dry and intact</li> <li>6. Child/ caregiver teaching</li> </ol>
PHYSICAL EXAM	<ul style="list-style-type: none"> <li>• Obtain history</li> <li>• Complete physical exam</li> <li>• Assess vital signs</li> <li>• Complete pain assessment (refer to <a href="#">Pain Assessment Guidelines</a>)</li> <li>• Obtain accurate in and out</li> </ul>	<ul style="list-style-type: none"> <li>• Complete pain assessment every 4 hours</li> <li>• Ensure child has adequate pain control (refer to <a href="#">Pain Management Guidelines</a>)</li> <li>• Monitor vital signs as per Bedside Pews (refer to <a href="#">Sepsis Clinical Pathway</a>)</li> <li>• Obtain accurate in and out</li> <li>• Complete wound assessment</li> <li>• Complete abdominal assessment</li> </ul>	
DIET & IV FLUIDS	<ul style="list-style-type: none"> <li>• Ensure that patient is NPO</li> <li>• Administer D5W and 0.9 NaCl with 20mmol KC/L at maintenance</li> <li>• Bolus as indicated</li> <li>• Refer to <a href="#">Fluid and Electrolyte Guidelines</a></li> </ul>	<ul style="list-style-type: none"> <li>• Clear fluids to regular diet as tolerated</li> <li>• IV to maintenance; TKVO once adequate oral fluid intake</li> <li>• Bolus as indicated</li> <li>• Refer to <a href="#">Fluid and Electrolyte Guidelines</a></li> </ul>	
MEDICATION	<ul style="list-style-type: none"> <li>• Ceftriaxone/Metronidazole IV; if allergy then Clindamycin or Ciprofloxacin &amp; Metronidazole. Refer to the e-formulary</li> <li>• Pain medication as needed; morphine/acetaminophen/ NSAIDs. Refer to the <a href="#">e-formulary</a></li> </ul>	<ul style="list-style-type: none"> <li>• Acetaminophen as needed for pain/fever</li> <li>• Ketorolac or ibuprofen as needed for pain management</li> <li>• Morphine IV bolus PRN</li> </ul>	
ACTIVITY & EDUCATION	<ul style="list-style-type: none"> <li>• Activity: as tolerated</li> <li>• Consent for surgery</li> <li>• Pre-op procedures for child and caregiver</li> <li>• Review parental involvement in care (pre and post-operatively)</li> </ul>	<p><b>Diet:</b></p> <ul style="list-style-type: none"> <li>• Advance diet as tolerated</li> </ul> <p><b>Pain:</b></p> <ul style="list-style-type: none"> <li>• Acetaminophen and ibuprofen (if not contraindicated) for 48 hours then as needed</li> <li>• Review need for pain management</li> </ul> <p><b>Incision care:</b></p> <ul style="list-style-type: none"> <li>• Leave steri-strips until fall off on own or remove after 10 days</li> <li>• Once steri-strips removed, may wash incision gently with soap and water</li> </ul> <p><b>Signs and symptoms of wound infection:</b></p> <ul style="list-style-type: none"> <li>• Fever</li> <li>• Redness around incision</li> <li>• Drainage from incision</li> <li>• Increasing pain around incision</li> </ul> <p><b>Bathing:</b></p> <ul style="list-style-type: none"> <li>• May shower or bathe; 48 hours after surgery</li> </ul> <p><b>Activity:</b></p> <ul style="list-style-type: none"> <li>• Ambulate in hallway</li> <li>• May return to normal daily activities as patient feels able</li> </ul>	<p><b>When to call surgeon's office:</b></p> <ul style="list-style-type: none"> <li>• Wound infection</li> <li>• Vomiting</li> <li>• Fever</li> <li>• Pain</li> </ul> <p><b>Follow-up:</b></p> <ul style="list-style-type: none"> <li>• Confirm need for follow-up with Primary Surgeon</li> <li>• Family doctor/pediatrician in 1-2 weeks</li> </ul>

[PRINTABLE VERSION](#)

©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.

## Inpatient Perforated Appendicitis Care Pathway

	DAY OF ADMISSION	DAY # 1	DAY # 2 - # 3	DAY # 4	DAY # 5
GOALS	<ol style="list-style-type: none"> <li>1. Hydration maintained</li> <li>2. Adequate pain control</li> <li>3. Patient prepared for OR if surgical management required</li> <li>4. Child/family are advised of pre-op bath. Wipes to be used upon arrival. Refer to <a href="#">procedure document</a></li> </ol>	<ol style="list-style-type: none"> <li>1. Afebrile</li> <li>2. Adequate pain control</li> <li>3. Ambulating</li> <li>4. Able to tolerate clears (immediately post-op)</li> <li>5. Incision intact, no drainage; dry and intact</li> <li>6. If Nasogastric Tube present, advance from intermittent suction to straight drainage</li> </ol>	<ol style="list-style-type: none"> <li>1. Afebrile</li> <li>2. Adequate pain control</li> <li>3. Ambulating</li> <li>4. Able to tolerate regular diet</li> <li>5. Incision dry and intact</li> <li>6. If Nasogastric Tube present, advance from straight drainage to clamp and remove</li> </ol>	<ol style="list-style-type: none"> <li>1. Afebrile</li> <li>2. Adequate pain control</li> <li>3. Ambulating</li> <li>4. Able to tolerate regular diet</li> <li>5. Incision dry and intact</li> <li>6. Child and family understand discharge teaching</li> <li>7. Able to tolerate oral antibiotics</li> </ol>	
PHYSICAL EXAM	<ul style="list-style-type: none"> <li>• History &amp; Physical</li> <li>• Vital Signs</li> <li>• Height and Weight</li> <li>• Pain Assessment (focus on abdominal) every 4 hours</li> <li>• Accurate In &amp; Out</li> </ul>		<ul style="list-style-type: none"> <li>• Vital signs as per BPEWS (<a href="#">Refer to Sepsis Clinical Pathway</a>)</li> <li>• Pain assessment (focus on abdominal) every 4 hours</li> <li>• Adequate pain control</li> <li>• Accurate In &amp; Out</li> <li>• Wound assessment (remove surgical dressing, leave steri-strips)</li> <li>• Abdominal assessment</li> </ul>		
IV FLUIDS	<ul style="list-style-type: none"> <li>• D5W &amp; 0.9% NaCl with 20mmol KCl/L</li> <li>• Bolus as clinically indicated with 0.9% NS or Lactated Ringer's</li> <li>• Refer to <a href="#">Fluid and Electrolyte Guidelines</a></li> </ul>	<ul style="list-style-type: none"> <li>• D5W &amp; 0.9% NaCl with 20mmol KCl/L at maintenance</li> <li>• Bolus as clinically indicated with 0.9% NS or Lactated Ringer's</li> <li>• Refer to <a href="#">Fluid and Electrolyte Guidelines</a></li> </ul>	<ul style="list-style-type: none"> <li>• Maintenance until adequate oral fluid intake and then TKVO</li> </ul>		
DIET			<ul style="list-style-type: none"> <li>• Diet as tolerated</li> <li>• If NPO, ensure that child is receiving IV fluids with D5W</li> <li>• Assess need for PN therapy</li> </ul>		
MEDICATIONS	<p>Pre-operatively:</p> <ul style="list-style-type: none"> <li>• Pain medication as needed (morphine / acetaminophen)</li> <li>• Start Q24h dosing of IV ceftriaxone and metronidazole (Refer to <a href="#">e-formulary</a> for dosing)</li> </ul> <p>Post-operatively:</p> <ul style="list-style-type: none"> <li>• Acetaminophen as needed for pain/fever</li> <li>• Morphine IV as required</li> <li>• Please check with Primary Surgeon if NSAIDs can be prescribed (Ketorolac vs. ibuprofen).</li> <li>• Q24h dosing of IV ceftriaxone and metronidazole until child is tolerating oral diet and afebrile for 12 hours (Refer to Appendicitis Management Pathway Algorithm (Refer to <a href="#">e-formulary</a> for dosing)</li> </ul>	<p>Pain management:</p> <ul style="list-style-type: none"> <li>• If on morphine infusion, wean as tolerated</li> <li>• Acetaminophen every 4 to 6 hours for 48 hours then as needed for pain/fever</li> <li>• Ketorolac or ibuprofen every 6 hours for 48 hours</li> </ul> <p>Antibiotics:</p> <ul style="list-style-type: none"> <li>• Q24h dosing of IV ceftriaxone and metronidazole</li> <li>• Consider switching to oral antibiotics - Clavulin (amoxicillin/clavulanate) to complete an additional 7day course) when child is afebrile for a minimum of 12 hours, tolerating oral diet and maintaining oral hydration and abdominal pain is well managed/improving</li> <li>• Refer to <a href="#">e-formulary</a> for dosages</li> <li>• Refer to <a href="#">Appendicitis Algorithm</a></li> </ul>			
ACTIVITY	<ul style="list-style-type: none"> <li>• Ambulating</li> </ul>		<ul style="list-style-type: none"> <li>• Ambulating to chair daily</li> <li>• Progress to ambulating in hallway X 5</li> </ul>		
DIAGNOSTIC IMAGING		<ul style="list-style-type: none"> <li>• Consider an abdominal ultrasound to evaluate for drainable intra-abdominal collection if child is not improving or clinically worsens</li> <li>• If collection is found, refer to <a href="#">Appendicitis Algorithm</a> for child with drainable collection</li> </ul>			
FAMILY / CAREGIVER EDUCATION	<ul style="list-style-type: none"> <li>• Pre-op procedures for parent and child</li> <li>• Consent for surgery signed</li> </ul>	<ul style="list-style-type: none"> <li>• When diet will be started</li> <li>• Need for pain management</li> <li>• Need for mobilizing</li> <li>• Parental involvement in care</li> </ul>	<p>Incision care:</p> <ul style="list-style-type: none"> <li>• Leave steri-strips until fall off on own or remove after 10 days</li> <li>• Once steri-strips removed, may wash incision gently with soap and water</li> </ul> <p>Signs and symptoms of wound infection:</p> <ul style="list-style-type: none"> <li>• Fever</li> <li>• Redness around incision</li> <li>• Drainage from incision</li> <li>• Increasing pain around incision</li> </ul> <p>Bathing:</p> <ul style="list-style-type: none"> <li>• May shower or bathe; 48 hours after surgery</li> </ul> <p>Activity:</p> <ul style="list-style-type: none"> <li>• Ambulate in hallway at least 5 times</li> <li>• May return to normal daily activities as patient feels able</li> </ul>	<p>When to call surgeon's office:</p> <ul style="list-style-type: none"> <li>• Wound infection</li> <li>• Vomiting</li> <li>• Fever</li> <li>• Pain</li> </ul> <p>Follow-up:</p> <ul style="list-style-type: none"> <li>• Confirm need for follow-up with Primary Surgeon</li> <li>• Family doctor/pediatrician in 1-2 weeks</li> </ul>	

## [PRINTABLE VERSION](#)

### 4.0 Related Documents

[E-formulary](#)

[Sepsis Pathway](#)

[Pain Management Guidelines](#)

[Pain Assessment Guidelines](#)

[Fluid & Electrolyte Guidelines](#)

©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.

## 5.0 References

1. Acker, S. N., Hurst, A. L., Bensard, D. D., Schubert, A., Dewberry, L., Gonzales, D., Parker, S. K., Tong, S., & Partrick, D. A. (2016). Pediatric appendicitis and need for antibiotics at time of discharge: Does Route of Administration matter? *Journal of Pediatric Surgery*, 51(7), 1170–1173. <https://doi.org/10.1016/j.jpedsurg.2016.03.004>
2. Alloo, J., Gerstle, T., Shilyansky, J. & Ein, S. (2004). Appendicitis in children less than 3 years of age: a 28 year review. *Pediatric Surgery International*, 19, 777-779.
3. Arnold, M. R., Wormer, B. A., Kao, A. M., Klima, D. A., Colavita, P. D., Cosper, G. H., Heniford, B. T., & Schulman, A. M. (2018). Home intravenous versus oral antibiotics following appendectomy for perforated appendicitis in children: A randomized controlled trial. *Pediatric Surgery International*, 34(12), 1257–1268. <https://doi.org/10.1007/s00383-018-4343-0>
4. Berrios-Torres, S. I et al., (2017). Centers for Disease Control and Prevention Guideline for the Prevention of Surgical Site Infection, 2017. *JAMA Surgery*. doi:10.1001/jamasurg.2017.0904
5. Bratzler, D. & Houck, P. (2004) Antimicrobial Prophylaxis for Surgery: An Advisory Statement from the National Surgical Infection Prevention Project. *Clinical Infectious Diseases*, 38, 1706-1715.
6. Children's Hospital of Philadelphia: Appendicitis without Known GI Disease Clinical Pathway- Emergency
7. Children's Hospital of Philadelphia: Appendicitis Clinical Pathway- Inpatients
8. Cincinnati Children's Hospital: Emergency Appendectomy Clinical Pathway
9. Dreznik, Y. (2018). Dual versus Triple Antibiotics Regimen in Children with Perforated Acute Appendicitis. *European Journal of Pediatric Surgery*, 28(06), 491-494
10. Ein, S., Langer, J., & Daneman, A. (2005). Nonoperative management of pediatric ruptured appendix with inflammatory mass or abscess: presence of an appendicolith predicts recurrent appendicitis. *Journal of Pediatric Surgery*, 40, 1612-1615.
11. Fraser, J., Aguayo, P., Leys, C., Keckler, S., Newland, J., Sharp, S., Murphy, J., Snyder, C., Sharp, R., Andrews, W., Holcomb, G., Ostlie, D., St. Peter, S. (2010). A complete course of intravenous antibiotics vs a combination of intravenous and oral antibiotics for perforated appendicitis in children: a prospective, randomized trial. *Journal of Pediatric Surgery* 45, 1198-1202.
12. Gee, K., Ngo, S., Burkhalter, L., & Beres, A. L. (2018). Safety and feasibility of same-day discharge for uncomplicated appendicitis: A prospective cohort study. *Journal of Pediatric Surgery*, 53, 988-990.
13. Groter, R. R., Sarah-May, M. L., Groter-Stam, M. A.W., Eker, H. H., Bakx, R., Lee, J. H., & Heij, H. (2017). Systematic review of nonoperative versus operative treatment of uncomplicated appendicitis. *Journal of Pediatric Surgery*, 52, 1219-1227.
14. Inpatient and Surgical Care: Appendectomy for Ruptured Appendix with Abscess or Generalized Peritonitis Care Guideline
15. Lansdale, N., Fryer, S., Stockdale, M., Bancroft, J., Orr, J., Corbett, H., & Kenny, S. (2019). Prospective evaluation of a clinical response directed pathway for complicated appendicitis. *Journal of Pediatric Surgery*, 54, 272-275.
16. Lee SL, Stark R, Yaghoobian A, Shekherdimian S, Kaji A. Does age affect the outcomes and management of pediatric appendicitis? *J Pediatr Surg*. 2011;46(12):2342-2345. doi:10.1016/j.jpedsurg.2011.09.030
17. Lee, S., Islam, S., Cassidy, L., Abdullah, F., Arca, M. (2010). Antibiotics and appendicitis in the pediatric population: an American Pediatric Surgical Association Outcomes and Clinical Trials Committee Systematic Review. *Journal of Pediatric Surgery*, 45, 2181-2185.
18. Lou, T., Falk, G.A., Burnweit, C. A., Ramos, C., Knight, C., & Malvezzi. (2016). Early transition to oral antibiotics for treatment of perforated appendicitis in pediatric patients: Confirmation of the safety and efficacy of a growing national trend. *Journal of Pediatric Surgery*, 51, 903-907.
19. Matthaiou DK, Peppas G, Bliziotis IA, Falagas ME. Ciprofloxacin/metronidazole versus  $\beta$ -lactam-based treatment of intra-abdominal infections: a meta-analysis of comparative trials. *Int J Antimicrob Agents*. 2006. doi:10.1016/j.ijantimicag.2006.04.005
20. Meier, D., Guzzetta, P., Barber, L., & Seetharamaiah, R. (2003). Perforated appendicitis in children: Is there a best treatment?. *Journal of Pediatric Surgery*, 38(10), 1520-1524.
21. Newman, K., Ponsky, T., Kittle, K., Dyk, L., Thropp, C., Giesecker, K., Sills, M. & Gilbert, J. (2003). Appendicitis 2000: Variability in Practice, Outcomes, and Resource Utilization at Thirty Pediatric Hospitals. *Journal of Pediatric Surgery*, 38, 372-379.
22. Rice HE, Brown RL, Gollin G, et al. Results of a pilot trial comparing prolonged intravenous antibiotics with sequential intravenous/oral antibiotics for children with perforated appendicitis. *Arch Surg*. 2001. doi:10.1001/archsurg.136.12.1391
23. RNAO Best Practice Guideline: Assessment and Management of Pain (2007)
24. Saluja, S., Sun, T., Mao, J., Steigman, S., Oh, S., Yeo, H. L., Sedrakyan, A., & Merianos, D. (2018). Early versus late surgical management of complicated appendicitis in children: A statewide database analysis within one-year follow-up. *Journal of Pediatric Surgery*, 53,1339-1344
25. Solomkin, JS, et al. Diagnosis and management of complicated intra-abdominal infection in adults and children: guidelines by the Surgical Infection Society and the Infectious Diseases Society of America [2]. *Clin Infect Dis* 2010;50:133-164.
26. St. Peter SD, Tsao K, Spilde TL, et al. Single daily dosing ceftriaxone and metronidazole vs standard triple antibiotic regimen for perforated appendicitis in children: a prospective randomized trial. *J Pediatr Surg*. 2008. doi:10.1016/j.jpedsurg.2008.02.018
27. Sujka, J.A., Weaver, K.L., Sobrino, J.A. et al. (2019). Efficacy of oral antibiotics in children with post-operative abscess from perforated appendicitis. *Pediatric Surgery International*, 35, 329-333 <https://doi.org/10.1007/s00383-018-4424-0>
28. University Hospitals of Cleveland: Pediatric Appendicitis/Simple Laparoscopic/Open Technique Care Path
29. Vaos G, Dimopoulou A, Gkioka E, Zavras N. Immediate surgery or conservative treatment for complicated acute appendicitis in children? A meta-analysis. *Journal of Pediatric Surgery*. 2018.
30. Warner, B. W., Kulick, R. M., Stoops, M.M, Mehta, S., Stephan, M., & Kotagal, U.R. (1998). An Evidence-Based Clinical Pathway for

©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.

Acute Appendicitis Decreases Hospital Duration and Cost.

31. Zani, A., Hall, N. J., Rahman, A., Morini, F., Prato, A. P., Friedmacher, F., Koivusalo, A., Heurn, E., & Pierro, A. (2019). European Paediatric Surgeon's Association Survey on the Management of Pediatric Appendicitis. *European Journal of Pediatric Surgery*, 29(01), 053-061

## Guideline Group and Reviewers

### Guideline Group Membership:

1. Monping Chiang, NP, General and Thoracic Surgery
2. Dr. Joshua Ramjist, Fellow, General and Thoracic Surgery
3. Natasha Brownrigg NP General Surgery
4. Dr. Annie Fecteau, Surgeon, General and Thoracic Surgery

### Internal Reviewers:

1. Kealey Clarke, RN, Quality Leader- 5B General Surgery
2. Sabrina Boohan, Clinical Pharmacist- 5B General Surgery
3. Christine McGovern, Sr Clinical Manager- 5B General Surgery

### Attachments:

[appendicitis algorithm.pdf](#)

[Non perforated appy pathway.pdf](#)

[Perforated appy pathway.pdf](#)

©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.