

Hypoxic Ischemic Encephalopathy (HIE) Clinical Care Pathway

Version: 8

This is a CONTROLLED document for internal use only, valid only if accessed from the Policies and Procedures site.

Content

- 1.0 [Introduction](#)
- 2.0 [Target Patient Population](#)
- 3.0 [Therapeutic Hypothermia Criteria](#)
- 4.0 [Neurologic Assessment](#)
 - 4.1 [Modified Sarnat Score](#)
 - 4.2 [Thompson Score](#)
- 5.0 [Clinical and Neurophysiologic Monitoring during Therapeutic Hypothermia](#)
 - 5.1 [Laboratory Monitoring](#)
- 6.0 [Patient Considerations](#)
- 7.0 [Process for Management of HIE in NICU](#)
- 8.0 [References](#)
- 9.0 [Guideline Group and Reviewers](#)

1.0 Introduction

Hypoxic ischemic encephalopathy (HIE) is defined as an abnormal neurologic state that occurs during the neonatal period following a hypoxic-ischemic insult. There is evidence from both animal and human studies that therapeutic hypothermia provides neuroprotection by reducing the severity of brain injury and resulting in an improved neurological outcome (1–4). All infants with HIE should be assessed for eligibility to receive therapeutic hypothermia. Evidence suggests that hypothermia in neonates with moderate to severe HIE reduces the severity of brain injury and leads to improved neurological outcomes (3). Therapeutic hypothermia (TH) is considered the standard of care in newborns with HIE.

2.0 Target Patient Population

- The commencement of therapeutic hypothermia **within 6 hours of birth**, for eligible patients, is the desired target (1,3,5). Recent data showed that the time to reach target temperature (i.e., therapeutic hypothermia initiated before or after 4 hours of age) demonstrated no significant effect on mortality, short-term outcomes, brain injury on MRI, and neurodevelopmental impairment at 2 years of age in infants with HIE (6).

©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.

- All term or late-preterm infants \geq 35 weeks gestation with moderate or severe HIE should be considered for therapeutic hypothermia.
- Therapeutic hypothermia is not recommended for preterm infants and infants with mild HIE as there is currently no evidence that cooling is beneficial (3). Studies have shown that therapeutic hypothermia in preterm infants less than 35 weeks gestation was associated with adverse effects (7) and hypothermia is associated with increased mortality and morbidity in very preterm infants (8). Similarly, there is insufficient evidence to recommend therapeutic hypothermia for infants with mild HIE (3). The benefits of therapeutic hypothermia in preterm infants and infants with mild HIE may not outweigh the risks, therefore therapeutic hypothermia is currently not recommended for these infants until more evidence is available.
- In the presence of profound encephalopathy where the risk of death or adverse neurodevelopmental outcome is high, the responsible physician may choose not to offer hypothermia treatment if there is no plan to pursue aggressive treatment.
- Initiation of hypothermia does not preclude a decision to withdraw life-sustaining therapy.

3.0 Therapeutic Hypothermia Criteria

Inclusion Criteria: Infant should fulfill all 4 criteria:	
1.	Gestational age (GA) greater than or equal to 35 weeks.
2.	Less than 6 hours post-delivery.
3.	Evidence of intrapartum hypoxia is defined as: <ol style="list-style-type: none"> Cord or postnatal blood gas within one hour of birth with pH less than or equal to 7.00 OR base excess (BE) of greater than or equal to -16. If pH between 7.01 – 7.15 or BE -10 to -15.9 and Apgar score is 5 or less at 10 minutes OR need for continued ventilation or resuscitation at 10 minutes. If there is no blood gas available then must have evidence of an acute perinatal event, i.e., placental abruption, uterine rupture, maternal trauma or cardiopulmonary arrest, late or persistent variable decelerations in an encephalopathy newborn.
4.	Signs of moderate or severe encephalopathy are defined as presence of clinical seizures OR 3 or more of the items in the moderate or severe categories using the modified Sarnat score.

Exclusion Criteria	
1.	Preterm infants born less than 35 weeks gestation.
2.	Neonates with mild encephalopathy on Sarnat Scoring.
3.	Neonates with a weight less than 1.8 kg.
4.	Clinically significant refractory coagulopathy despite treatment.
5.	Moribund neonates, or neonates with major congenital or genetic abnormalities, in whom no further aggressive treatment is planned.

©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.

4.0 Neurological Assessment

Standardized neurological examination must be performed by a physician or nurse practitioner skilled in neurological assessment within the first hour of life to determine the degree of encephalopathy. The gold standard in the encephalopathy assessment is the modified Sarnat Score; however, it is recommended to also perform the Thompson score as an additional tool to trend clinical evolution numerically. Of note, cerebral function monitoring using amplitude-integrated EEG (aEEG) can assist in screening infants for eligibility for therapeutic hypothermia, although this should not be used to exclude otherwise eligible neonates as per Newborn Brain Society Guidelines (9).

4.1 Modified Sarnat Score

	Normal	Mild	Moderate	Severe
Level of consciousness	<input type="checkbox"/> Alert and responsive	<input type="checkbox"/> Hyperalert, jittery, exaggerated responses	<input type="checkbox"/> Lethargic	<input type="checkbox"/> Stupor or coma
Spontaneous activity	<input type="checkbox"/> Normal	<input type="checkbox"/> Decreased ± periods of hyperactivity	<input type="checkbox"/> Decreased	<input type="checkbox"/> No activity
Posture	<input type="checkbox"/> Predominantly flexed posture	<input type="checkbox"/> Mild distal flexion	<input type="checkbox"/> Strong distal flexion or complete extension	<input type="checkbox"/> Intermittently decerebrate
Tonus	<input type="checkbox"/> Flexor tone in extremities	<input type="checkbox"/> Slightly increased peripheral tone	<input type="checkbox"/> Hypotonia or hypertonia	<input type="checkbox"/> Flaccid or rigid
Primitive reflexes 1) Suck 2) Moro	<input type="checkbox"/> Strong <input type="checkbox"/> Strong	<input type="checkbox"/> Weak, poor <input type="checkbox"/> Low threshold to elicit	<input type="checkbox"/> Weak or bites only <input type="checkbox"/> Incomplete	<input type="checkbox"/> Absent <input type="checkbox"/> Absent
Autonomic function 1) Pupils 2) Heart Rate 3) Respirations	<input type="checkbox"/> Normal size, reactive <input type="checkbox"/> Normal heart rate <input type="checkbox"/> Normal	<input type="checkbox"/> Mydriasis <input type="checkbox"/> Tachycardia (>160/min) <input type="checkbox"/> Hyperventilation	<input type="checkbox"/> Miosis <input type="checkbox"/> Bradycardia (<100/min) <input type="checkbox"/> Periodic breathing	<input type="checkbox"/> Skewed/Non-reactive <input type="checkbox"/> Variable <input type="checkbox"/> Apnea/On ventilator

4.2 Thompson Score

Score	0	1	2	3
Tone	Normal	Hypertonic	Hypotonic	Flaccid
Level of consciousness	Normal	Hyper alert stare	Lethargic	Comatose
Seizures	None	Infrequent < 3/day	Frequent > 2/day	
Posture	Normal	Fisting, cycling	Distal flexion	Decerebrate
Moro	Normal	Partial	Absent	
Grasp	Normal	Poor	Absent	
Suck	Normal	Poor	Absent ± bites	
Respiration	Normal	Hyperventilation	Brief apnea	IPPV (apnea)
Fontanel	Normal	Full, not tense	Tense	
TOTAL SCORE				

©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.

5.0 Clinical and Neurophysiologic Monitoring during Therapeutic Hypothermia

- The modified Sarnat score and Thompson score should be administered daily and documented in the Electronic Health Record (EHR) in all newborns undergoing TH until the end of rewarming.
- Newborns receiving therapeutic hypothermia should receive aEEG monitoring during the cooling and rewarming periods.
- Continuous EEG should be considered if there is a concern for seizures or suppressed background patterns On aEEG (discontinuous normal voltage, burst-suppression, continuous low voltage, isoelectric trace) (10).
- Near-infrared spectroscopy (NIRS) should be connected to newborns undergoing therapeutic hypothermia to assess brain hemodynamics (11). Refer to [Near-Infrared Spectrometry \(NIRS\) Use in the NICU](#).
- A medical order for specific alarm limits is to be placed for patients requiring alarm limits to be set outside the standard parameters.

5.1 Laboratory Monitoring

The following **laboratory monitoring** is recommended as a minimum and is to be ordered by the medical team. Additional laboratory tests may be ordered as needed.

Timing	Lab Testing
POCT glucose	Every 6 hours throughout the cooling process
On admission	Gas, lactate, CBC, coagulation, electrolytes, ALT, AST, urea, creatinine, ammonia, calcium, glucose
12 hours after initiation of cooling	Gas, lactate, electrolytes, C-reactive protein
24 hours after initiation of cooling	- Gas, lactate, CBC, coagulation, electrolytes, LFTs, bilirubin, urea, creatinine, ammonia, calcium, phosphate. - Ensure newborn screen (NBS) is collected.
48 hours after initiation of cooling	Electrolytes, urea and creatinine, bilirubin
72 hours after initiation of cooling	Glucose, electrolytes, bilirubin, CBC
24 hours after rewarming has been completed	Glucose, electrolytes, calcium, bilirubin

Infants Without a Clear Sentinel Event

- For infants presenting without a clear sentinel event, other causes of neonatal encephalopathy should also be considered and investigated appropriately.
- The possibility of a systemic and/or central nervous system infection should be ruled out.
- Placental pathology should be sent, and pathology results should be monitored closely.
- Neurometabolic disorders are a frequent cause of neonatal encephalopathy, so a metabolic/genetic consultation for further testing with whole exome sequencing is strongly advised in these infants in consultation with the neonatal neurology team.

©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.

6.0 Patient Considerations

- **Venous and Arterial Access**
 - Consider venous and arterial access needs for patient monitoring prior to the cooling process as difficulties in obtaining access may occur due to hypothermia-associated decreased perfusion.
- **Fluid and Nutritional Requirements**
 - Fluid and nutritional requirements should be assessed daily and when there are changes to the level of patient sedation. Hypothermia, sedation, and the effects of a hypoxic ischemic insult have an additive effect on the infant's metabolic activity.
- **Enteral Feeding during Therapeutic Hypothermia**
 - Patients with moderate-severe HIE will remain NPO during cooling.
 - Patients with milder HIE may be eligible for trophic feeds – refer to [NICU Enteral Feeding and Gastroesophageal Reflux Management Guideline](#) for [Enteral Feeding During Therapeutic Hypothermia](#).
- **Minimizing the Risk of Subcutaneous Fat Necrosis**
 - Once target temperature has been reached **do not use additional ice packs** on the skin as this increases the risk for subcutaneous fat necrosis. Refer to [Subcutaneous Fat Necrosis](#) guideline.
- **Analgesia and Sedation Management**
 - Indications for sedation include agitation or a shivering response.
 - Shivering leads to increased peripheral muscle oxygenation consumption. Neonates with excessive shivering should be considered for treatment with dexmedetomidine as per [e-formulary](#) (12) and refer to [NICU Sedation Guidelines](#).
 - Analgesia may be used instead of a sedative in the case of patients who have pain due to trauma at delivery, with low dose morphine infusion. Infants with hypoxic ischemic encephalopathy have reduced morphine clearance and elevated serum morphine concentrations.
 - Potentially toxic serum concentrations of morphine may occur with moderate hypothermia and infusion rates greater than 10 micrograms/kg per hour (13).
- **Concurrent Use of Anti-Seizure Medications (ASM) and Sedative Medications**
 - Caution should be used in patients who are receiving anticonvulsants in addition to sedative or opioid agents.
 - Please discontinue sedatives for patients receiving phenobarbital or midazolam infusions for seizures.
 - Refer to the [SickKids Neonatal Seizure Management Guidelines](#) for the management of seizures and consult Neonatal Neurology service for further guidance.

▪ **Holding during Therapeutic Hypothermia**

- Patients MAY be eligible for holding during cooling. Refer to [Infant Holding in the NICU](#) guideline for eligibility and procedure.

▪ **Rewarming**

- Rewarming should occur no faster than 0.5 degrees Celsius per hour, even if rewarming is occurring for other clinical indications such as hemodynamic instability or persistent coagulopathy. Refer to [Thermal Regulating System Use in NICU](#) for rewarming procedure.

7.0 Process for Management of HIE in NICU

1. Refer to the [Hypoxic Ischemic Encephalopathy \(HIE\) Care Pathway](#).
2. Refer to [Thermal Regulating System Use in the NICU \(i.e. Blanketrol®\)](#) guideline for how to provide therapeutic hypothermia.

8.0 References

1. Gluckman PD, Wyatt JS, Azzopardi D, Ballard R, Edwards AD, Ferriero DM, Polin RA, Robertson CM, Thoresen M, Whitelaw A, Gunn AJ. Selective head cooling with mild systemic hypothermia after neonatal encephalopathy: multicentre randomised trial. *Lancet*. 2005 Feb;365(9460):663–70.
2. Jacobs SE, Berg M, Hunt R, Tarnow-Mordi WO, Inder TE, Davis PG. *Cochrane Database Syst Rev*. 2013. Cooling for newborns with hypoxic ischaemic encephalopathy.
3. Lemyre B, Chau V. Hypothermia for newborns with hypoxic-ischemic encephalopathy. *Paediatr Child Health*. 2018 Jul;23(4):285–91.
4. Shankaran S, Laptook AR, Ehrenkranz RA, Tyson JE, McDonald SA, Donovan EF, Fanaroff AA, Poole WK, Wright LL, Higgins RD, Finan NN, Carlo WA, Duara S, Oh W, Cotten CM, Stevenson DK, Stoll BJ, Lemons JA, Guillet R, Jobe AH; Whole-body hypothermia for neonates with hypoxic-ischemic encephalopathy. *N Engl J Med*. 2005 Oct;353(15):1574–84.
5. Shah D, Mackay M, Lavery S. Accuracy of bedside electroencephalographic monitoring in comparison with simultaneous continuous conventional electroencephalography for seizure detection in term infants. *Pediatrics*. 121(6):1146–54.
6. Rao R, Comstock BA, Wu TW, Mietzsch U, Mayock DE, Gonzalez FF, Wood TR, Heagerty PJ, Juul SE, Wu YW. Time to Reaching Target Cooling Temperature and 2-year Outcomes in Infants with Hypoxic-Ischemic Encephalopathy. *J Pediatr*. 2024 Mar;226(113853).
7. Roca-LLabrés P, Fontalvo-Acosta M, Aldecoa-Bilbao V, Alarcón A. Therapeutic hypothermia in preterm infants under 36 weeks: Case series on outcomes and brain MRI findings. *Eur J Pediatr*. 2025 Jan;184(1):113.
8. Hogeveen M, Hooft L, Onland W. Adverse Outcomes in Very Preterm Infants: A Systematic Review. *Pediatrics*. 2025 May;155(5).

©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.

9. El-Dib M, Abend NS, Austin T, Boylan G, Chock V, Cilio MR, et al. Neuromonitoring in neonatal critical care part I: neonatal encephalopathy and neonates with possible seizures. *Pediatr Res.* 2023 Jul;94(1):64–73.
10. Chock VY, Rao A, Van Meurs KP. Optimal neuromonitoring techniques in neonates with hypoxic ischemic encephalopathy. *Front Pediatr.* 2023 Mar 8;11(1138062).
11. Mitra S, Bale G, Meek J, Tachtsidis I, Robertson NJ. Cerebral Near Infrared Spectroscopy Monitoring in Term Infants With Hypoxic Ischemic Encephalopathy-A Systematic Review. *Front Neurol.* 2020 May 27;11:393.
12. Baserga M, DuPont TL, Ostrander B, Minton S, Sheffield M, Balch AH, Bahr TM, Watt KM. Dexmedetomidine Use in Infants Undergoing Cooling Due to Neonatal Encephalopathy (DICE Trial): A Randomized Controlled Trial: Background, Aims and Study Protocol. *Front Pain Res.* 2021 Dec 7;2(770511).
13. Roka A, Melinda KT, Vasarhelyi B, Machay T, Azzopardi D, Szabo M. Elevated morphine concentrations in neonates treated with morphine and prolonged hypothermia for hypoxic ischemic encephalopathy. *Pediatrics.* 2008;121(4):e844–9.

9.0 Guideline Group and Reviewers

- The guideline was reviewed and updated December 2025.
- This guideline was critically reviewed and revised by SickKids Neonatal Neurocritical Care Team in May 2023.
- **Lead Authors:** Mehmet N. Cizmeci and Rhandi Christensen
- **Guideline Update Group Members:** Amr El-Shahed, Diane Wilson
- **Critical Review by Neonatal Neurocritical Care Team Members:** Brian Kalish, Cecil Hahn, Emily Tam, Linh Ly, Vann Chau, Vanna Kazazian. In addition, review completed by NICU Quality Management
- **Internal Reviewers:** Christine Elliott, Christopher Tomlinson

©The Hospital for Sick Children ("SickKids"). All Rights Reserved. This document was developed solely for use at SickKids. SickKids accepts no responsibility for use of this material by any person or organization not associated with SickKids. A printed copy of this document may not reflect the current, electronic version on the SickKids Intranet. Use of this document in any setting must be subject to the professional judgment of the user. No part of the document should be used for publication without prior written consent of SickKids.